

5-2018

# Effects of Oxalate Oxidase Transgene Expression on Mycorrhizal Colonization in American Chestnuts

Hannah Roden

Follow this and additional works at: <http://digitalcommons.esf.edu/honors>

 Part of the [Genetics and Genomics Commons](#), and the [Plant Breeding and Genetics Commons](#)

---

## Recommended Citation

Roden, Hannah, "Effects of Oxalate Oxidase Transgene Expression on Mycorrhizal Colonization in American Chestnuts" (2018).  
*Honors Theses*. 125.  
<http://digitalcommons.esf.edu/honors/125>

This Thesis is brought to you for free and open access by Digital Commons @ ESF. It has been accepted for inclusion in Honors Theses by an authorized administrator of Digital Commons @ ESF. For more information, please contact [digitalcommons@esf.edu](mailto:digitalcommons@esf.edu), [cjkoons@esf.edu](mailto:cjkoons@esf.edu).

Effects of oxalate oxidase transgene expression on mycorrhizal colonization in American Chestnuts

By

Hannah Roden  
Candidate for Bachelor of Science  
Environmental and Forest Biology  
With Honors

May 2018

**APPROVED**

Thesis Project Advisor: \_\_\_\_\_  
Thomas R. Horton, Ph.D.

Second Reader: \_\_\_\_\_  
William M. Powell, Ph.D.

Honors Director: \_\_\_\_\_  
William M. Shields, Ph.D.

Date: \_\_\_\_\_

## Abstract

American chestnuts have been genetically engineered to express an oxalate oxidase gene isolated from wheat, which allows the plants to degrade oxalic acid, the virulence factor produced by *Cryphonectria parasitica*, the pathogenic fungus that causes the chestnut blight. In this study, we investigated the effect of oxalate oxidase gene expression on rate of mycorrhizal colonization of American chestnut root tips. Six-month-old transgenic and wildtype plantlets were placed in a soil inoculant and cultured in a greenhouse for 7 months, then root tips were visually assessed to estimate how many were colonized (expressed as a percentage). Three genotypes were used: ELLIS 1 trees were a cloned wildtype, and Darling 54 and Darling 58 were ELLIS 1 genotypes transformed with the oxalate oxidase gene, differing only in where the gene construct was inserted into the genome. Of surviving trees, 100% of both ELLIS 1 and Darling 58 trees were >95% colonized. Ninety percent of Darling 54 were >95% colonized, and 10% were 90-95% colonized. A Fisher's exact test of independence was used to reach the conclusion that expression of the oxalate oxidase gene has no effect on mycorrhizal colonization, a result consistent with similar studies.

## Table of Contents

Acknowledgements.....	i
Thesis Essay.....	1
<i>Introduction</i> .....	1
<i>Methods</i> .....	2
<i>Results</i> .....	4
<i>Discussion</i> .....	4
Sources Cited and Consulted.....	7
Appendices.....	10

## **Acknowledgements**

I am grateful for the help of Dr. Horton and Dr. Powell, who designed and setup the study, gave advice and proofread this draft. I thank Dr. Horton for providing equipment and lab space, as well instruction in the lab. He taught me important lessons in sampling and lab procedure. I also thank Tyler Desmarais for his help and advice, and his support in the greenhouse. Tyler oversaw the care of the plants from tissue culture, inoculation, and throughout the course of mycorrhizal colonization, by maintaining a water and pesticide regimen.

## Thesis essay

### *Introduction*

American chestnuts (*Castanea dentata*) have been genetically transformed to express an oxalate oxidase gene isolated from wheat (*Triticum aestivum*; Lane et al. 1993). Oxalate oxidase is an enzyme that breaks down oxalic acid, the virulence factor produced by *Cryphonectria parasitica* (Chen et al. 2010), the chestnut blight fungus, to necrotize the cambium tissue of non-resistant American chestnut trees. By degrading oxalic acid, American chestnut trees become resistant to the disease (B. Zhang et al. 2013), not by killing the fungus or preventing infection, but by preventing the lethal effects and making the fungus non-virulent. Resistant trees do not harm the fungus.

The oxalate oxidase gene used to produce disease-resistant American chestnuts was isolated from wheat, but it is a gene also found in other monocots (Lane et al. 1993; Dahiya et al. 2010), so while it is a foreign gene to the chestnut, it is native to the ecosystem. However, before transformed trees are introduced to the wild for restoration, it is imperative to measure non-target environmental effects. In this study, a greenhouse bioassay was conducted, and chestnut roots were observed to measure the potential effect of the expression of oxalate oxidase on mycorrhizal colonization.

The mycorrhizal network is used by plants to enhance water and mineral absorption. Fungi are able to access these nutrients because they have a unique enzymatic capacity (Qian, Wang and Yin, 2012). They also increase the plant's root volume and can access soil pores smaller than those plant roots can reach (Dighton, 2009). Study has shown that American chestnut seedlings form ectomycorrhizal associations.

Ectomycorrhizal fungi do not penetrate the cell wall of the host plant, instead forming an intercellular interface (Dulmer, LeDuc, and Horton, 2015; Palmer, Lindner, and Volk, 2008). The effects of oxalate oxidase expression on mycorrhizal colonization of American chestnut have been previously studied and with no effect on ectomycorrhizal colonization rates (D'Amico et al. 2015). The influence of various disease resistance-enhancing transgenes in multiple plants species on soil microbe communities have also been studied, with similar results (Becker et al. 2014; Cheeke et al. 2015; Kaldorf et al. 2002; Kaur et al. 2017; Newhouse et al. 2007; Vierheilig et al. 1995; Xie et al. 2016; Zeng et al. 2014; Zhang et al. 2015). A few studies showed reduced arbuscular mycorrhizal infection in transformed Bt cotton (Beura et al. 2015; Chen et al. 2016). It is expected that we will observe no significant difference in mycorrhizal colonization between wildtype and transformed American chestnuts.

In this study, the rate of mycorrhizal colonization was compared between three American chestnut genotypes. The first was ELLIS1, a cloned wildtype. The others were Darling54 and Darling58, ELLIS1 genotypes transformed with the oxalate oxidase gene, differing only in the genome insertion sites of the oxalate oxidase (OxO) gene construct. These transgenic trees differ from ones in previous studies (D'Amico et al. 2015) by fewer genome inserts, fewer selectable marker genes present, and higher OxO gene expression levels.

### *Methods*

Soil samples for mycorrhizal inoculum were collected from ESF's Lafayette Road Experiment Station in Syracuse, New York, in a mixed hardwood stand. 23 random soil samples were dried, then sifted through a 0.5cm mesh. Samples were dried to select for

resistant propagules such as fungal spores and sclerotia. The soil inoculant was made by combining a ratio of 1:1:2 dried soil to sand to sphagnum peat moss. The inoculant was split evenly among 45 cylindrical pots, which had been previously sterilized overnight in a 7% bleach solution. About 6-month old tissue culture-generated *C. dentata* were bare-rooted and transplanted into the pots containing the inoculant. Three types of *C. dentata* were used, 15 individuals (n=15) each of 'Ellis 1' 'Darling 54' and 'Darling 58' (**Table 1**). Trees were cultured in a greenhouse until colonization took place, about 5 months. Greenhouse temperatures ranged from 18 to 29 degrees Celsius. Plants received a 16-hour photo period under high pressure sodium lights emitting light equivalent to full sun in field conditions. Plants were watered with tap water every two or three days. Tap water had a generally high pH and low electrical conductivity, or salt/fertilizer content.

Mycorrhizal colonization rate was assessed by collecting a continuous root length of at least 15cm after six months in the greenhouse. All root tips along the sample were observed, and the percentage of root tips that were colonized (categorized as mycorrhizal), and uncolonized (categorized as nonmycorrhizal) were visually estimated and assigned to categorical percentage ranks (less than 5%, more than 95%, or within ranges of 10%). A root tip was considered mycorrhizal if it was actively colonized or senescent with indications of previous colonization. The frequencies of each category in each treatment were calculated. A Fisher's exact test of independence, with a significance of 0.05, was used to test the null hypothesis that there was no significant difference in colonization between any two treatments.



## *Results*

Mortality rate was 0% among ELLIS1 genotypes, 33% among Darling54 genotypes, and 20% among Darling58 genotypes. All surviving trees had greater than 90% mycorrhizal colonization. 100% of root tips were colonized in all Darling 58 and Ellis1 trees, and 90% of Darling54 trees. 90-95% of root tips were colonized in 10% of Darling54 (**Table 1**).

## *Discussion*

There were no statistical differences between the rates of mycorrhizal colonization of the three genotypes of American chestnut, two of which were engineered to express oxalate oxidase. As previously stated, American chestnut associates with ectomycorrhizal fungi, which take part in chemical exchange with the plant host without breaking the plant cell wall or plasma membrane. Oxalate oxidase produced as a result of the expression of the transgene may not come into contact with the mycorrhizal fungi if it is present in the cytoplasm, apoplast, or some other intercellular structure and is not exported past the cell wall, or if it is differentially expressed in tissues and not root tissues.

If the gene is expressed in the roots, it may have no effect on mycorrhizae because oxalic acid production is associated with pathogens and saprobes, specifically brown-rot fungi that rely on the breakdown of plant tissues for survival (Green and Clausen, 2003; Hastrup et al. 2012; Na et al. 2017). Mycorrhizae depend on a symbiosis with their plant hosts, an exchange of water and minerals for the carbohydrate products of

photosynthesis, and do not rely on the action of oxalic acid, so its absence would likely have little effect on mycorrhizal colonization.

Hydrogen peroxide and carbon dioxide are the byproducts of oxalic acid decomposition. Hydrogen peroxide is an oxidative agent that is sometimes toxic to some mycelial fungi in high concentrations (El-Gazzar and Marth, 1988; Zharare, Kabanda and Poku, 2010), but it had no negative effects on the arbuscular mycorrhizae that colonized our chestnuts. This may be because hydrogen peroxide is associated with the temporal and spatial control of arbuscular mycorrhizae colonization; hydrogen peroxide was observed to accumulate in cortical root cells occupied by arbuscules, the haustorial organs of arbuscular mycorrhizae, and around hyphal tips attempting to penetrate host tissues (Salzer, Corbière and Boller, 1999). It has also been found that arbuscular mycorrhizae use hydrogen peroxide as a signal molecule in a pathway that enhances plant host resistance to pathogens (R. Zhang et al. 2013). Increasing the concentration of hydrogen peroxide, to a certain degree, would not harm the mycorrhizae, as they create and use this compound themselves. However, it may be harmful in very high amounts, and future study should look at hydrogen peroxide accumulation in transgenic American chestnut tissues over time, as oxalic acid is continuously broken down.

The results of this experiment are comparable to the results of similar studies (Becker et al. 2014; Cheeke et al. 2015; Kaldorf et al. 2002; Kaur et al. 2017; Newhouse et al. 2007; Vierheilig et al. 1995; Xie et al. 2016; Zeng et al. 2014; Zhang et al. 2015). One such study also investigated the mycorrhizal colonization of wildtype and blight-resistant genetically engineered American chestnut, as well as related species including red oak, American beech, and Chinese chestnut, and found that mycorrhizal colonization

rates differed between soil types, but not between tree genotypes (D'Amico et al. 2015). In another study, wheat was transformed with an apoplast-targeted defensin gene which granted the wheat significant resistance to infection by fungal rust. However, expression of the foreign gene had no effect on the plant's symbiotic relationship with mycorrhizal fungi (Kaur et al., 2016). A third study found that vesicular-arbuscular mycorrhizal colonization was not affected by various constitutively-expressed pathogenesis-related proteins, including chitinases, in tobacco plants (Vierheilig, 1995). In a final study, fungal symbionts were not affected by transgenic cotton expressing a foreign insecticidal gene that targeted lepidopteran pests. There were differences between the population size and structure of rhizosphere soil fungi between the transformed and conventional cotton only at some stages of plant growth. But fungal population size and structure were significantly influenced by variation in years and the stage of plant growth, and there was no difference in dominant fungi between cotton genotypes at any stage, indicating the transgenic, pest-resistant cotton had no significant effect on fungal community size or structure (Xie et al. 2016). These studies corroborate our results, demonstrating that expression of transgenes which enhance disease resistance have little or no effect on mycorrhizal communities.

## Sources cited and consulted

- Becker, Regina, Ben Bubner, Rainer Remus, Stephan Wirth, and Andreas Ulrich. 2014. "Impact of Multi-Resistant Transgenic Bt Maize on Straw Decomposition and the Involved Microbial Communities." *Applied Soil Ecology* 73 (January):9–18. <https://doi.org/10.1016/j.apsoil.2013.08.002>.
- Beura, Kasturikasen, Amit Kumar Pradhan, Amitava Rakshit, and Mahendra Singh. 2015. "Effect of Bt Transgene on AM Fungi Infection and Yield Attributes of Cotton (*Gossypium Hirsutum* L.)." *International Journal of Bio-Resource & Stress Management* 6 (4):517–22. <https://doi.org/10.5958/0976-4038.2015.00077.9>.
- Cheeke, Tanya E., Ursel M. Schütte, Chris M. Hemmerich, Mitchell B. Cruzan, Todd N. Rosenstiel, and James D. Bever. 2015. "Spatial Soil Heterogeneity Has a Greater Effect on Symbiotic Arbuscular Mycorrhizal Fungal Communities and Plant Growth than Genetic Modification with *Bacillus Thuringiensis* Toxin Genes." *Molecular Ecology* 24 (10):2580–93. <https://doi.org/10.1111/mec.13178>.
- Chen, C., Sun, Q., Narayanan, B., Nuss, D. L., & Herzberg, O. (2010). Structure of Oxalacetate Acetylhydrolase, a Virulence Factor of the Chestnut Blight Fungus. *Journal of Biological Chemistry*, 285 (34), 26685–26696
- Dahiya, T., Yadav, S., Chauhan, N., Handa, P., & Pundir, C. S. (2010). Strawberry Fruit Oxalate Oxidase — Detection, Purification, Characterization and Physiological Role. *Journal of Plant Biochemistry and Biotechnology*, 19(2), 247–250. <https://doi.org/10.1007/BF03263349>
- D'Amico, Katherine M., Thomas R. Horton, Charles A. Maynard, Stephen V. Stehman, Allison D. Oakes, and William A. Powell. 2015. "Comparisons of Ectomycorrhizal Colonization of Transgenic American Chestnut with Those of the Wild Type, a Conventionally Bred Hybrid, and Related Fagaceae Species." *Applied and Environmental Microbiology* 81 (1):100–108. <https://doi.org/10.1128/AEM.02169-14>.
- Dighton, J. 2009. "Mycorrhizae." *Encyclopedia of Microbiology (Third Edition)*: 153-162. <https://doi-org.esf.idm.oclc.org/10.1016/B978-012373944-5.00327-8>
- Dulmer, Kristopher M., Stephen D. LeDuc, and Thomas R. Horton. 2014. "Ectomycorrhizal Inoculum Potential of Northeastern US Forest Soils for American Chestnut Restoration: Results from Field and Laboratory Bioassays." *Mycorrhiza* 24 (1):65–74. <https://doi.org/10.1007/s00572-013-0514-y>.
- El-Gazzar, Fathy E. and Elmer H. Marth. 1988. "Role of Hydrogen Peroxide in the Prevention of Growth and Aflatoxin Production by *Aspergillus parasiticus*." *Journal of Food Protection* 51 (4):263-168.
- Green III, Frederick, Carol A. Clausen. 2003. "Copper tolerance of brown-rot fungi: time course of oxalic acid production." *International Biodeterioration & Biodegradation* 51 (2):145-149 [https://doi.org/10.1016/S0964-8305\(02\)00099-9](https://doi.org/10.1016/S0964-8305(02)00099-9)

- Hastrup, Anne C.S., Frederick Green III, Patricia K. Lebow, Bo Jensen. 2012. "Enzymatic oxalic acid regulation correlated with wood degradation in four brown-rot fungi." *International Biodeterioration & Biodegradation* 75:109-114 <https://doi.org/10.1016/j.ibiod.2012.05.030>
- Kaldorf, Michael, Matthias Fladung, Hans-J. Muhs, and François Buscot. 2002. "Mycorrhizal Colonization of Transgenic Aspen in a Field Trial." *Planta* 214 (4):653–60. <https://doi.org/10.1007/s004250100658>.
- Kaur, Jagdeep, John Fellers, Alok Adholeya, Siva L. S. Velivelli, Kaoutar El-Mounadi, Natalya Nersesian, Thomas Clemente, and Dilip Shah. 2017. "Expression of 'Apoplast-Targeted Plant Defensin <Emphasis Type='Italic'>MtDef4.2</Emphasis> Confers Resistance to Leaf Rust Pathogen <Emphasis Type='Italic'>Puccinia Triticina</Emphasis> but Does Not Affect Mycorrhizal Symbiosis in Transgenic Wheat." *Transgenic Research* 26 (1):37–49. <https://doi.org/10.1007/s11248-016-9978-9>.
- Lane BG, Dunwell JM, Ray JA, Schmitt MR, Cuming AC. 1993. "Germin, a protein marker of early plant development, is an oxalate oxidase." *J Biol Chem.* 1993;268:12239–12242
- Na, Ren, Yanjie Luo, Haoyu Bo, Jian Zhang, Ruifang Jia, Qinglin Meng, Hongyou Zhou, Jianjun Hao, Jun Zhao. 2017. "Responses of sunflower induced by *Sclerotinia Sclerotiorum* infection." *Physiological and Molecular Plant Pathology.* <https://doi.org/10.1016/j.pmpp.2017.12.004>
- Newhouse, Andrew E., Franziska Schrodt, Haiying Liang, Charles A. Maynard, and William A. Powell. 2007. "Transgenic American Elm Shows Reduced Dutch Elm Disease Symptoms and Normal Mycorrhizal Colonization." *Plant Cell Reports* 26 (7):977–87. <https://doi.org/10.1007/s00299-007-0313-z>.
- Palmer, Jonathan M., Daniel L. Lindner, and Thomas J. Volk. 2008. "Ectomycorrhizal Characterization of an American Chestnut <Emphasis Type='SmallCaps'>(</Emphasis> <Emphasis Type='Italic'>Castanea Dentata</Emphasis> <Emphasis Type='SmallCaps'>)-Dominated Community in Western Wisconsin." *Mycorrhiza* 19 (1):27. <https://doi.org/10.1007/s00572-008-0200-7>.
- Qian, Kulmai, Liping Wang and Ningning Yin. 2012. "Effects of AMF on soil enzyme activity and carbon sequestration capacity in reclaimed mine soil." *International Journal of Mining Science and Technology* 22 (4):553-557
- Salzer, Peter, Hélène Corbiere, Thomas Boller. 1999. "Hydrogen peroxide accumulation in *Medicago truncatula* roots colonized by the arbuscular mycorrhiza-forming fungus *Glomus intraradices*." *Planta* 208 (3):319-325. <https://doi.org/10.1007/s004250050565>
- Vierheilig, H., M. Alt, J. Lange, M. Gut-Rella, A. Wiemken, and T. Boller. 1995. "Colonization of Transgenic Tobacco Constitutively Expressing Pathogenesis-Related Proteins by the Vesicular-Arbuscular Mycorrhizal Fungus *Glomus Mosseae*." *Applied and Environmental Microbiology* 61 (8):3031–34.

Xie, Ming, Yan-Jun Zhang, De-Liang Peng, Gang Wu, Peng Xu, Jin-Jin Zhao, and Zhao-Rong Zhang. 2016. "Field Studies Show No Significant Effect of a Cry1Ab/Ac Producing Transgenic Cotton on the Fungal Community Structure in Rhizosphere Soil." *European Journal of Soil Biology* 73 (March):69–76. <https://doi.org/10.1016/j.ejsobi.2016.01.006>.

Zeng, Huilan, Fengxiao Tan, Yanyan Zhang, Yuanjiao Feng, Yinghua Shu, and Jianwu Wang. 2014. "Effects of Cultivation and Return of *Bacillus Thuringiensis* (Bt) Maize on the Diversity of the Arbuscular Mycorrhizal Community in Soils and Roots of Subsequently Cultivated Conventional Maize." *Soil Biology and Biochemistry* 75 (August):254–63. <https://doi.org/10.1016/j.soilbio.2014.04.024>.

Zhang, B., Oakes, A. D., Newhouse, A. E., Baier, K. M., Maynard, C. A., & Powell, W. A. 2013. A threshold level of oxalate oxidase transgene expression reduces *Cryphonectria parasitica* induced necrosis in a transgenic American chestnut (*Castanea dentata*) leaf bioassay. *Transgenic Research*, 22(5), 973–982. <https://doi.org/10.1007/s11248-013-9708-5>

Zhang, Rui-Qin, Hong-Hui Zhu, Hai-Quan Zhao, and Qing Yao. 2013. "Arbuscular mycorrhizal fungal inoculation increases phenolic synthesis in clover roots via hydrogen peroxide, salicylic acid and nitric oxide signaling pathways." *Journal of Plant Physiology* 170 (1):74-79. <https://doi.org/10.1016/j.jplph.2012.08.022>

Zhang, Yan-Jun, Ming Xie, Gang Wu, De-Liang Peng, and Wen-Bin Yu. 2015. "A 3-Year Field Investigation of Impacts of Monsanto's Transgenic Bt-Cotton NC 33B on Rhizosphere Microbial Communities in Northern China." *Applied Soil Ecology* (May):18–24. <https://doi.org/10.1016/j.apsoil.2015.01.003>.

Zharare, Godfrey E., Siti M. Kabanda and Judith Z. Poku. 2010. "Effects of temperature and hydrogen peroxide on mycelial growth of eight *Pleurotus* strains." *Scientia Horticulturae* 125 (2):95-102. <https://doi.org/10.1016/j.scienta.2010.03.006>

## Appendices

**Table 1.** Tree mortality, as a percentage of the sample size (n=15) and colonization of ectomycorrhizal fungi of wildtype and transgenic American chestnut (*C. dentata*) plantlets, measured in % root tips colonized.

<u>Tree Type</u>	<u>Number of surviving trees observed</u>	<u>Tree mortality</u>	<u>Surviving trees with &gt;95% mycorrhizal colonization</u>	<u>Percent of surviving trees with 90-95% mycorrhizal colonization</u>	<u>Percentage of surviving trees with &lt;90% mycorrhizal colonization</u>
‘Ellis 1’	15	None	15 (100%)	None	None
‘Darling 54’	10	5 (33%)	9 (90%)	1 (10%)	None
‘Darling 58’	12	3 (20%)	12 (100%)	None	None